



NRL/MR/7430--04-8842

Compilation of Data Sources Used to Construct Bottom-Sediment Province Maps for the United States East Coast Continental Shelf from Cape Kennedy to Cape Hatteras and for Various Indonesian Waters

W. CHAD VAUGHAN

*Seafloor Sciences Branch
Marine Geosciences Division*

FREDERICK A. BOWLES

*Omni Technologies, Inc.
New Orleans, LA*

JOSHUA W. CHATELAIN

*Department of Geography
University of New Orleans
New Orleans, LA*

December 27, 2004

20050214 048

Approved for public release; distribution is unlimited.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 27-12-2004		2. REPORT TYPE Memorandum Report		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Compilation of Data Sources Used to Construct Bottom-Sediment Province Maps for the United States East Coast Continental Shelf from Cape Kennedy to Cape Hatteras and for Various Indonesian Waters				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) W. Chad Vaughan, Frederick A. Bowles,* and Joshua W. Chatelain†				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Marine Geosciences Division Stennis Space Center, MS 39529-5004				8. PERFORMING ORGANIZATION REPORT NUMBER NRL/MR/7430-04-8842	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Oceanographic Office Stennis Space Center, MS 39529				10. SPONSOR / MONITOR'S ACRONYM(S) NAVO	
				11. SPONSOR / MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES *Omni Technologies, Inc., New Orleans, LA †Department of Geography, University of New Orleans, New Orleans, LA					
14. ABSTRACT The purpose of this report is to document the data sources used to construct maps and digital databases of bottom-sediment type for a portion of the continental shelf off the eastern United States and for Indonesian waters. The east coast mapping region extends from approximately 28 deg-30 min N (Cape Canaveral) to 35 deg-30 min N (Cape Hatteras). Mapping of the Indonesian waters includes the Banda, Ceram, Molucca, Halmahera, and Flores Seas (see the attached figures showing the approximate boundaries of the mapped areas).					
15. SUBJECT TERMS Sediment; Bottom; Maps; United States; Indonesia; Digital; Database					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UL	18. NUMBER OF PAGES 19	19a. NAME OF RESPONSIBLE PERSON W. Chad Vaughan
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code) 228-688-5369

CONTENTS

INTRODUCTION	1
APPROACH	1
EAST COAST, UNITED STATES	2
NIMA Charts	2
References (Journals, Reports)	2
Other Data Sources	5
References of Marginal or No Use	6
Potential Data Sources (Not Acquired)	7
INDONESIAN WATERS	7
NIMA Charts Used	7
References Used	7
Land Geology References (Used to Predict Near-shore Sediments)	9
References of Marginal or No Use	9

Compilation of Data Sources used to Construct Bottom-Sediment Province Maps for the United States East Coast Continental Shelf from Cape Kennedy to Cape Hatteras and for Various Indonesian Waters

Introduction

The purpose of this report is to document the data sources used to construct maps and digital databases of bottom-sediment type for a portion of the continental shelf off the eastern United States and for Indonesian waters. The east coast mapping region extends from approximately 28 deg-30 min N (Cape Canaveral) to 35 deg-30 min N (Cape Hatteras). Mapping of the Indonesian waters includes the Banda, Ceram, Molucca, Halmahera, and Flores Seas (see the attached figures showing the approximate boundaries of the mapped areas).

Approach

The following is a brief outline of the approach used to construct the bottom-sediment maps for each region. A more detailed discussion can be found in Bowles and Phelps (1999; Naval Research Laboratory Report NRL/MR/7432—99-8223).

1. National Imagery and Mapping Agency (NIMA) nautical charts are used as the working (base) charts for each region. These charts are readily available from the NAVOCEANO library and feature bottom sediment data in addition to bathymetric data and coastlines.
2. An extensive literature and institutional search is performed to identify and acquire existing bottom-sediment data in each region. These data are in the form of point data (core, dredge, grab samples) with associated sediment-type descriptors, or existing sediment-type boundary maps.
3. As data is acquired, it is carefully assessed, geo-referenced (in the case of existing sediment maps) and transferred to the base charts. Once all the data is transferred, boundaries are drawn around the point data designating areas (provinces) of similar sediment type. These province boundaries are then merged with the boundaries from existing sediment maps.
4. Upon completion of the maps, the information on each map is digitized. The digitized information consists of separate files for land boundaries, sediment-province boundaries, and point data. The point

data files consist of individual files for sand, silt, clay, rock, etc. The sediment nomenclature used to identify bottom type follows the enhanced "200 Categories" set defined in the current NAVOCEANO Master Sediment Tables.

EAST COAST, UNITED STATES

NIMA Charts

11480	11510	11523
11488	11511	11531
11490	11512	11532
11502	11513	11535
11503	11516	11536
11504	11517	11537
11506	11520	11539
11508	11521	11542
11509	11522	11543

References (journals, reports)

Ayers, M., Blackwelder, B. W., Howard, J. D., Keer, F., Knebel, H., and Pilkey, O. H., 1980. Vibracore studies: Georgia Embayment shelf. Final report; environmental studies, southeastern United State Atlantic outer continental shelf 1977; geology. U. S. Geological Survey Open-File Report 6.1-6.90.

Batten, R. W., 1961. The sediments of the Beaufort Inlet area, North Carolina. Southeastern Geology, 3(1), 191-205.

Cazeau, C. J., Du Bar, J. R., and Henry, J. S., 1963. Notes on marine geology off the mouth of the North Edisto River, South Carolina. Southeastern Geology, 5(1), 157-176.

Cleary, W. J. and Pilkey, O. H., 1968, Sedimentation in Onslow Bay. Southeastern Geology, 9, 1-17.

- Fluke, L. A., 1994. Recent Atlantic shelf sedimentation within a siliciclastic-carbonate transition, Florida, USA. M.S. Thesis, Florida Institute of Technology, Melbourne.
- Goodell, H. G., 1967. The sediments and sedimentary geochemistry of the southeastern Atlantic shelf. *Journal of Geology*, 75(6), 665-692.
- Grosz, A. E. and Nelson, D. D., 1989. Textural and mineralogical analyses of surficial sediments offshore of Myrtle Beach, South Carolina. United States Geological Survey Open-File Report 89-169, Reston, VA.,
- Henry, V. J. and Hoyt, J. H., 1968. Quaternary paralic and shelf sediments of Georgia. *Southeastern Geology*, 9, 195-214.
- Hollister, C. D., 1973. Atlantic continental shelf and slope of the United States – texture of surface sediment from New Jersey to Southern Florida. U.S. Geological Survey Professional Paper 529-M, 23p.
- Irving, C. S. Jr. and Siegel, F. R., 1969. Distribution and provenance of minerals from continental shelf sediments off the South Carolina coast. *Journal of Sedimentary Petrology*, 39(1), 276-296.
- Kirby-Smith, W. W., 1989. The community of small macroinvertebrates associated with rock outcrops on the continental shelf of North Carolina. National Oceanographic and Atmospheric Administration-National Undersea Research Program Report 89-2.
- Macintyre, I. C. and Pilkey, O. H., 1969. Preliminary comments on linear sand-surface features, Onslow Bay, North Carolina continental shelf: problems in making detailed sea-floor observations. *Maritime Sediments*, 5(1), 26-29.
- Marinelli, R. L., Jahnke, R., A., Craven, D. B., Nelson, J. R., and Eckman, J. E., 1998. Sediment nutrient dynamics on the South Atlantic Bight continental shelf. *Limnology and Oceanography*, 43(6), 1305-1320.
- McNich, J. E. and Wells, J. T., 1999. Sedimentary processes and depositional history of a cape-associated shoal, Cape Lookout, North Carolina. *Marine Geology*, 158, 233-252.

Meisburger, E. P., 1977. Sand resources on the inner continental shelf of the Cape Fear region, North Carolina. Miscellaneous Report No. 77-11, U.S. Army, Corps of Engineers, Coastal Engineering Research Center, Fort Belvoir, VA., 20p.

Meisburger, E. P., 1979. Reconnaissance geology of the inner continental shelf, Cape Fear region, North Carolina. Technical Paper No. 79-3, U. S. Army, Corps of Engineers, Coastal Engineering Research Center, Fort Belvoir, VA., 135p.

Meisburger, E. P. and Field, M. E., 1975. Geomorphology, shallow structure, and sediments of the Florida inner continental shelf, Cape Canaveral to Georgia. Technical Memorandum No. 54, U. S. Army, Corps of Engineers, Coastal Engineering Research Center, Fort Belvoir, VA.. 119p.

Meisburger, E. P. and Field, M. E., 1976. Neogene sediments of Atlantic inner continental shelf off northern Florida. American Association of Petroleum Geologists Bulletin, 60(11), 2019-2037.

Milliman J. D., Pilkey, O. H., and Blackwelder, B. W., 1968. Carbonate sediments on the continental shelf, Cape Hatteras to Cape Romain. Southeastern Geology, 9, 245-267.

Milliman, J. D., 1972. Atlantic continental shelf and slope of the United States – petrology of the sand fraction of the sediments, northern New Jersey to southern Florida: U.S. Geological Survey Professional Paper 529-J, 40p.

Pickett, T. E. and Ingram, R. L., 1969. The modern sediments of Pamlico Sound, North Carolina. Southeastern Geology, 11(2), 53-

Pilkey, O. H. and Terlecky, P. M., 1966. Distribution of surface sediments on the Georgia continental shelf. In: Hoyt, J. H., Henry, V. J. Jr., and Howard, J. D., Pleistocene and Holocene sediments, Sapelo Island, Georgia, and vicinity, Geological Society of America Southeastern Section, Field Trip No. 1, 28-39.

Pilkey, O. H. and Luternauer, J. L., 1967. A North Carolina shelf phosphate deposit of possible commercial interest. Southeastern Geology, 9(1), 33-51.

Pilkey, O. H., Keer, F., and Keer, S., 1980. Surficial sediments of the U.S. Atlantic southeastern United States Atlantic continental shelf. Final report; environmental studies, southeastern United State Atlantic outer continental shelf 1977; geology. U. S. Geological Survey Open-File Report 5.1-5.40.

Pinet, P. R. and Frey, R. W., 1977. Organic carbon in surface sands seaward of Altamaha and Doboy Sounds, Georgia. Geological Society of America Bulletin, 88, 1731-1739

Richardson, M. D. and Briggs, K. B., 1993. On the use of acoustic impedance values to determine sediment properties. Proceedings of the Institute of Acoustics, 15, Part 2, 15-24.

Schmitter, R. D. and Freeman-Lynde, R. P., 1988. Horblende content of Georgia-South Carolina, U.S.A., nearshore sands: support for shoreward sediment transport in the nearshore zone. Sedimentary Geology, 57, 153-169.

Thieler, E. R., Brill, A. L., Cleary, W. J., Hobbs III, C. H., and Gammisch, R. A., 1995. Geology of the Wrightsville Beach, North Carolina shoreface: implications for the concept of shoreface profile of equilibrium. Marine Geology, 126, 271-287.

Other Data Sources

Duke University. Unpublished sediment maps with grain size and/or percent carbonate data for core sites along the east coast of the U. S. (furnished by Orrin Pilkey).

Naval Oceanographic Office, Lab Item 813 (Sediment Laboratory).

Naval Research Laboratory. Diver-collected samples from Onslow Bay during Operation Unified Spirit (USS Champion).

U.S. Geological Survey Continental Margin (CONMAR) Sediment file. (Downloaded from National Geophysical Data Center website)

World Data Center for Marine Geology and Geophysics. Index to Marine Geological Samples Database. National Geophysical Data Center, Boulder, CO.

References of Marginal or No Use

- Bucca, P. J. and Sellinger, C., 1985. A preassessment of environmental conditions on the continental shelf adjacent to Jacksonville, Florida. Naval Ocean Research and Development Activity (NORDA) Report 88, Stennis Space Center, MS, 39529, 23p.
- Gorsline, D., 1963. Bottom sediments of the Atlantic shelf and slope off the southern United States. *Journal of Geology*, 71, 422-440.
- Milliman, J. D., Pilkey, O. H., and Ross, D. A., 1972. Sediments of the continental margin off the eastern United States. *Geological Society of America Bulletin*, 83, 1315-1334.
- Pilkey, O. H. and Frankenberg, D., 1964. The relict-recent sediment boundary on the Georgia continental shelf. *Georgia Academy of Science Bulletin*, 22(1), 37-40.
- Reed, T., L., 1976. Heavy metal concentrations in the sediments of a portion of the east Florida continental shelf. M.S. Thesis, Florida Institute of Technology, Melbourne (covers an area south of this mapping project).
- Riggs, S. R., Snyder, S. W., Hine, A. C., and Mearns, D. L., 1996. Hardbottom morphology and relationship to the geologic framework: mid-Atlantic continental shelf. *Journal of Sedimentary Research*, 66(4), 830-846.
- Riggs, S. R., Ambrose Jr., W. G., Cook, J. W., Snyder, S. W., and Snyder, S. W., 1998. Sediment production on sediment-starved continental margins: the interrelationship between hardbottoms, sedimentological and benthic community processes, and storm dynamics. *Journal of sedimentary Research*, 68(1), 155-168.
- Riggs, S. R., Snyder, S. W. P., Hine, A. C., Snyder, S. W., Ellington, M. D., and Mallette, P. M., 1985. Geologic framework of phosphate resources in Onslow Bay, North Carolina continental shelf. *Economic Geology*, 80, 716-738.

Potential Data Sources (Not Acquired)

Howard, J. D., 1974. Surficial sediments of the Georgia continental shelf. Georgia Geological Survey Bulletin, 87, 135p.

Kingery, F. A., 1973. Textural analysis of shelf sands off the Georgia coast. Unpublished M.S. Thesis, California State University, San Diego.

Manheim, F. T. and Herring, J. R., 1989. Location and core background data. In: Phosphorite potential in the continental shelf off Georgia; results of the TACTS core studies. Open-File Report – U. S. Geological Survey, 4-24.

Rochna, D. A., 1961. Physiography and sedimentology of the "Outer Banks" between Cape Lookout and Ocracoke Inlet, N. C. Unpublished M.S. Thesis, University of Kansas.

INDONESIAN WATERS:

NIMA Charts Used

72131	73018	73291
73000	73020	73451
73002	73022	73461
73004	73024	73470
73006	73034	73471 (2981)
73008	73130	73481
73010	73141	73760
73012	73261	
73016	73271	

References Used

Anonymous, 1985. Report on marine meology. Geological Research and Development Centre (Indonesia). Committee for Co-ordination of Joint Prospecting for Mineral Resources in Asian Offshore Areas (CCOP). Proceedings of the 12th Session, 192-198.

Budiono, K. and Kuntoro, 1994. Coastal and marine geological studies for development of the Kendari coastal area, south east Sulawesi, Indonesia.

Committee for Co-ordination of Joint Prospecting for Mineral Resources in Asian Offshore Areas (CCOP). Proceedings of the 29th Annual Session, 299-313.

Dwiyanto, B., Soeprapto, T. A. and Budiono, K., 1992. Marine geology and geophysics of Ambon Bay, Molluca-Indonesia. *Jurnal Geologi dan Sumberdaya Mineral (Journal of Geology and Mineral Resources)*, 2(12), 2-16.

Ganssen, G., Troelstra, S. R., Faber, B., van der Kaars, W. A. and Situmorang, M., 1989. Late Quaternary palaeoceanography of the Banda Sea, eastern Indonesia piston cores (Snellius-II Expedition, Cruise G5). *Netherlands Journal of Sea Research*, 24(4), 491-494.

Nitzsche, M., 1989. Submarine slope instability, eastern Banda Sea. *Netherlands Journal of Sea Research*, 24(4), 431-436.

Situmorang, M., 1989. Lithofacies and depositional pattern of sea floor sediments in the North Banda Sea, Indonesia. *Netherlands Journal of Sea Research*, 24(4), 405-413.

Situmorang, M. and Sarmili, L., 1997. Composition, morphology, dispersal patterns of gravel clasts and basement rocks in the Banda Arc sea floor, Eastern Indonesia. *Bulletin of the Marine Geology*, 12(1), 1-26.

Van Bergen Henegouw, C. N. and Richards, A. F., 1989. Geotechnical property variability over an active plate margin in the eastern Banda Sea. *Netherlands Journal of Sea Research*, 24(4), 437-444.

Van Hinte, J. E. and Shipboard Party, 1986. The Snellius-II Expedition. Progress Report. Theme I. Geology and Geophysics of the Banda Arc and Adjacent Areas; Cruise G5: Flores, Timor, Banda and Arafura Sea. Royal Netherlands Academy of Arts and Sciences-Indonesian Institute of Sciences, 79p.

Van der Linde W. J. M. and Shipboard Party, 1986. The Snellius-II Expedition. Progress Report. Theme I. Geology and Geophysics of the Banda Arc and Adjacent Areas; Cruise G5: Banda, Seram, Halmahera and

Maluku Sea. Royal Netherlands Academy of Arts and Sciences-Indonesian Institute of Sciences, 38p.

Land Geology References (used to predict near-shore sediments)

Charlton, T. R., Hall, R. and Partoyo, E., 1991. The geology and tectonic evolution of Waigeo Island, NE Indonesia. *Journal of Southeast Asian Earth Sciences* 6(3/4) 289-297.

Mugroto, B. and Ali, J., 1998. Tectonic rotations indicated by the Late Cenozoic paleomagnetic rocks in Buton Island, southeast Sulawesi, Indonesia. Coordinating Committee for Coastal and Offshore Geoscience Programmes in East and Southeast Asia (CCOP). Proceeding of the 33th Session, 246-258.

Nichols, G. J. and Hall, R., 1991. Basin formation and Neogene sedimentation in a backarc setting, Halmahera, eastern Indonesia. *Marine and Petroleum Geology*, 8, 50-61.

Ryacudu, R., Tossin, S. and Purnomo, E., 1998. Ampana strike-slip fault and its significance for hydrocarbon entrapment in the eastern arm of Sulawesi, Indonesia. Coordinating Committee for Coastal and Offshore Geoscience Programmes in East and Southeast Asia (CCOP). Proceedings of the 34th Session, 1-21.

Sufni Hakim, A. and Hall, R., 1991. Tertiary volcanic rocks from the Halmahera Arc, eastern Indonesia. *Journal of Southeast Asian Earth Sciences*, 6(3/4), 271-287.

Surono, 1998. Sedimentology of the oolitic limestone succession of the Paleogene Tampakura Formation, southeastern Sulawesi, Indonesia. Coordinating Committee for Coastal and Offshore Geoscience Programmes in East and Southeast Asia (CCOP). Proceeding of the 33th Session. 10-41.

References of Marginal or No Use

Adisaputra, M. K., 1989. Planktonic foraminifera in Recent bottom sediments of the Flores, Lombok, and Savu Basins, eastern Indonesia. *Netherlands Journal of Sea Research*, 24(4), 465-475.

Barmawidjaja, D. M., de Jong, A. F. M., van der Borg, K., van der Kaars, W. A. and Zachariasse, W. J., 1989. Kau Bay, Halmahera, a Late Quaternary palaeoenvironmental record of a poorly ventilated basin. *Netherlands Journal of Sea Research*, 24(4), 591-605.

Gullentops, F. and Monty, C., 1988. Sedimentology and petrology of Quaternary sediments in the Weber, Seram and Aru troughs, Indonesia. (Abstract) International Association of Sedimentologists, 9th European regional meeting: Leuven, Belgium, Sept. 1988, p.186.

Marle, L. J. van, Hinte, J. E. van and Nederbragt, A. J., 1987. Plankton percentage of the foraminiferal fauna in seafloor samples from the Australian-Irian Jaya continental margin, eastern Indonesia. *Marine Geology*, 77, 151-156.

Middelburg, J. J., Calvert, S. E. and Karlin, R., 1991. Organic-rich transitional facies in silled basins: response to sea-level change. *Geology*, 19(7), 679-682

Rehault, J. P., Malod, J. A., Larue, M., Burhanuddin, S. and Sarmili, L., 1991. A new sketch of the central North Banda Sea, Eastern Indonesia. *Journal of Southeast Asian Earth Sciences* 6(3/4) 329-334.

Van der Kaars, W. A., 1989. Aspects of Late Quaternary, palynology of eastern Indonesian deep-sea cores. *Netherlands Journal of Sea Research*, 24(4), 495-500..

Van der Linden, W. J. M., Hartosukohardjo, S. and Sukardjoho, H., 1989. Kau Bay, Halmahera: Regional setting, physiography and shallow structure. *Netherlands Journal of Sea Research*, 24(4), 573-581.

Van Paverd, P. J. and Bjorklund, K. R., 1989. Frequency distribution of polycystines radiolarians in surface sediments of the Banda Sea, eastern Indonesia. *Netherlands Journal of Sea Research*, 24(4), 511-521.

Van Waveren, I. M., 1989. Palynofacies analysis of surface sediments from the northeastern Banda Sea (Indonesia). *Netherlands Journal of Sea Research*, 24(4), 501-509.

Villeneuve, M., Cornee, J.-J., Martini, R., Zaninetti, J.-P., Burthanudin, S., and Malod, J., 1994. Upper Triassic shallow water limestones in the Sinta Ridge (Banda Sea, Indonesia). *Geo-Marine Letters* 14, 29-35.

Vroon, P. Z., van Bergen, M. J., Klaver, G. J. and White, W. M., 1995. Strontium, neodymium, and lead isotopic and trace-element signatures of the east Indonesian sediments : provenance and implications for Banda Arc magma genesis. *Geochimica et Cosmochimica Acta*, 59(12), 2573-2598.









